

NIKHEF/2006-009

HIGHER DIMENSIONAL VSI SPACETIMES AND SUPERGRAVITY

ANDREA FUSTER

*NIKHEF, Kruislaan 409, 1098 SJ Amsterdam, The Netherlands
fuster@nikhef.nl*

NICOS PELAVAS

*Department of Mathematics and Statistics, Dalhousie University, Halifax, Nova Scotia B3H
3J5, Canada
pelavas@mathstat.dal.ca*

We present the explicit form of higher dimensional VSI spacetimes in arbitrary number of dimensions. We discuss briefly the VSI's in the context of supergravity/strings.

1. Introduction

We are interested in higher dimensional spacetimes for which curvature invariants of all orders vanish (VSI spacetimes), contained within this class are the higher dimensional pp-waves. In general, the higher dimensional VSI spacetimes have Ricci and Weyl type III.¹ However, it is desirable to obtain explicit metric functions as has been done in four dimensions.² We present metrics for Ricci type N, Weyl type III VSI spacetimes.

2. Higher dimensional VSI spacetimes

Any N -dimensional VSI metric can be written in the form^{3,4}

$$ds^2 = 2du [dv + H(u, v, x^k) du + W_i(u, v, x^k) dx^i] + dx^i dx^i \quad (1)$$

where u, v are light-cone coordinates and $x^i, i = 1, \dots, N-2$, are real spatial coordinates. The functions H, W_i are real-valued. Note that (1) is a subclass of the higher dimensional Kundt metrics.³ Restricting (1) to be of Ricci type N results in the Einstein equation $R_{uu} = \Phi$, where Φ is determined by the matter field (see Appendix). Two distinct cases arise depending on whether the functions W_i depend on the light-cone coordinate v ($\epsilon = 1$) or not ($\epsilon = 0$):

$$W_1 = -\frac{2\epsilon}{x^1} v + W_1^{(0)}(u, x^k), \quad W_j = W_j^{(0)}(u, x^k)$$

Here $j = 2, \dots, N-2$; the superscript (0) indicates functions without v -dependence. The case $\epsilon = W_i^{(0)} = 0$ corresponds to higher dimensional pp-waves. It is perhaps

surprising that only one of the W_i functions is allowed to depend on v , as in the 4-D case. The resulting spacetimes are summarized in Table 1. An important remark is that the function $W_1^{(0)}$ can be gauged away by a coordinate transformation, the corresponding results have been recently presented.⁵ In this note we treat all $W_i^{(0)}$ functions equally, thereby revealing additional algebraic symmetries found in the metric functions of solutions. This complements the paper by Coley et al, wherein further details and explanation may be found.

ϵ	Weyl	Metric functions	Eq.
0	III	$W_i = W_i^{(0)}(u, x^k)$ $H = H^{(0)}(u, x^i) + \frac{1}{2} \left(F - W^{(0)i}{}_{,i} \right) v$, $F(u, x^i)$ defined by $F_{,i} = \Delta W_i^{(0)}$	A.1
	N	$W_i = x^k B_{ki}(u)$, $B_{ki}(u)$ antisymmetric $H = H^{(0)}(u, x^i)$	A.2
	O	W_i as in type N $H^{(0)} = \frac{1}{2} W^i W_i + x^i f_i(u) + H_\Phi^{(0)}$	A.2 A.3
1	III	$W_1 = -\frac{2}{x^1} v + W_1^{(0)}(u, x^k)$ $W_j = W_j^{(0)}(u, x^k)$ $H = H^{(0)}(u, x^i) + \frac{1}{2} \left(\tilde{F} - W^{(0)i}{}_{,i} - \frac{2W_1^{(0)}}{x^1} \right) v + \frac{v^2}{2(x^1)^2}$, $\tilde{F}(u, x^i)$ defined by $\tilde{F}_{,1} = \frac{2}{x^1} W^{(0)i}{}_{,i} + \Delta W_1^{(0)}$, $\tilde{F}_{,j} = \Delta W_j^{(0)}$	A.4
	N	$W_1 = -2\frac{v}{x^1} + x^j B_{j1}(u) + C_1(u)$ $W_j = x^i B_{ij}(u) + C_j(u)$, $B_{j1}(u), B_{ij}(u)$ antisymmetric $H = H^{(0)}(u, x^i) - \frac{W_1^{(0)}}{x^1} v + \frac{v^2}{2(x^1)^2}$	A.5
	O	W_i as in type N $H^{(0)} = \frac{1}{2} (W_1^{(0)})^2 + \frac{1}{2} \sum_j (W_j - x^1 B_{1j})^2 + x^1 g_0(u) + x^1 x^i g_i(u) - \frac{1}{16} \Phi_0(u) x^1 x^i x_i$, $\Phi = \Phi_0(u) x^1$	A.5

Table 1: All higher dimensional VSI spacetimes of Ricci type N. Ricci type O (vacuum) for $\Phi = 0$ in (A.1)-(A.5).

3. Outlook: VSI's in supergravity

We can consider the embedding of VSI spacetimes in supergravity. The idea is to construct bosonic solutions of the supergravity equations of motion where the metric is that of a VSI.⁶⁻⁸ This has been extensively studied in the literature in the case of (Weyl type N) pp-waves. These are exact string solutions to all orders in α' . The corresponding proof relies on the pp-waves invariants property. Furthermore, supergravity pp wave solutions preserve supersymmetry. The VSI metrics presented here generalize the pp-waves while the invariants property is maintained. Therefore it is natural to expect VSI supergravity solutions to have some of the pp-wave solutions' features. In fact, a certain class of VSI supergravity solutions has already been proved to be exact to all orders in α' .⁷ Supersymmetry properties have only been studied in the case of VSI's with a covariantly constant null vector.⁸ In conclusion, these results provide an opportunity to explore VSI supergravity solutions, and their associated supersymmetries.

Appendix A. Einstein equations

In the equations below $W_{ik} = W_{i,k}^{(0)} - W_{k,i}^{(0)}$ and $H^{(1)}$ is the coefficient of v in the metric function H .

$$\Delta H^{(0)} - \frac{1}{4} W_{ik} W^{ik} - 2H^{(1)}{}_{,i} W^{(0)i} - H^{(1)} W^{(0)i}{}_{,i} - W^{(0)i}{}_{,iu} + \Phi = 0 \quad (\text{A.1})$$

$$\Delta H^{(0)} - 2 \sum_{i < j} B_{ij}^2 + \Phi = 0 \quad (\text{A.2}) \quad H_{\Phi,ii}^0 = -\frac{1}{8} \Phi, \quad H_{\Phi,ik}^0 = 0 \quad (\text{A.3})$$

$$x^1 \Delta \left(\frac{H^{(0)}}{x^1} \right) + \left(\frac{W^{(0)j} W_j^{(0)} - (W_1^{(0)})^2}{x^1} \right)_{,1} + \frac{2}{x^1} (W_1^{(0)} W_{1,1}^{(0)} - W^{(0)j} W_{1,j}^{(0)}) \\ - 2H^{(1)}{}_{,i} W^{(0)i} - H^{(1)} W^{(0)i}{}_{,i} - \frac{1}{4} W_{ik} W^{ik} - W^{(0)i}{}_{,iu} + \Phi = 0 \quad (\text{A.4})$$

$$x^1 \Delta \left(\frac{H^{(0)}}{x^1} \right) - \frac{1}{(x^1)^2} \left((W_1^{(0)})^2 + \sum_j (W_j - x^1 B_{1j})^2 \right) - \sum_j B_{1j}^2 - 2 \sum_{i < j} B_{ij}^2 + \Phi = 0 \quad (\text{A.5})$$

References

1. A. Coley, R. Milson, V. Pravda and A. Pravdova, *Class. Q. Grav.* **21**, 5519 (2004).
2. V. Pravda, A. Pravdova, A. Coley and R. Milson, *Class. Q. Grav.* **19**, 6213 (2002).
3. A. Coley, R. Milson, N. Pelavas, V. Pravda, A. Pravdova and R. Zalaletdinov, *Phys. Rev. D* **67**, 104020 (2003).
4. A. Coley, S. Hervik and N. Pelavas, *Class. Q. Grav.* **23**, 3053 (2006).
5. A. Coley, A. Fuster, S. Hervik and N. Pelavas, *Class. Q. Grav.* **23**, 7431 (2006).
6. V. Frolov and F. Lin, *Phys. Rev. D* **73**, 104028 (2006).
7. A. Coley, *Phys. Rev. Lett.* **89**, 281601 (2002).
8. E. Bergshoeff, R. Kallosh and T. Ortin, *Phys. Rev. D* **47**, 5444 (1993).